

EQUIVARIANT EULER CHARACTERISTICS OF DISCRIMINANTS OF REFLECTION GROUPS

GRAHAM DENHAM AND NICOLE LEMIRE

ABSTRACT. Let G be a finite, complex reflection group acting on a complex vector space V , and δ its discriminant polynomial. The fibres of δ admit commuting actions of G and a cyclic group. The virtual $G \times C_m$ character given by the Euler characteristic of a fibre is a refinement of the zeta function of the geometric monodromy, calculated in [8]. We show that this virtual character is unchanged by replacing δ by a slightly more general class of polynomials. We compute it explicitly, by studying the poset of normalizers of centralizers of regular elements in G , and the subspace arrangement given by the proper eigenspaces of elements of G . As a consequence we also compute orbifold Euler characteristics and find some new “case-free” information about the discriminant.

1. SUMMARY

Let G be a finite reflection group acting on the vector space $V = \mathbf{C}^\ell$. Let \mathcal{A} denote the set of reflecting hyperplanes of G . For each $H \in \mathcal{A}$, let $\alpha_H \in V^*$ be a linear functional with kernel H . The discriminant polynomial δ of G is defined to be

$$\delta = \prod_{H \in \mathcal{A}} \alpha_H^{e_H},$$

where e_H is the order of the subgroup of G that fixes H pointwise. δ is the G -invariant polynomial of smallest degree whose zero set is exactly the set of reflecting hyperplanes. Let $m = \deg \delta$.

The fibres of δ over \mathbf{C}^* are diffeomorphic, by a theorem of Milnor [15]; let $F = \delta^{-1}(1)$, the Milnor fibre of δ . The action of G on V restricts to an action on F . At the same time, a cyclic group C_m acts on F , generated by a geometric monodromy map $h : F \rightarrow F$ defined by $h(x) = e^{2\pi i/m}x$.

The actions of G and C_m commute. Let $\Gamma = G \times C_m$. Then $H_\bullet(F, \mathbf{C})$ is a finite-dimensional representation of Γ . In this paper, we consider the Euler characteristic of F , valued in the character ring of Γ . That is, we define a virtual character χ_Γ by

$$(1.1) \quad \chi_\Gamma(F)(g) = \sum_{p \geq 0} (-1)^p \text{Tr}(g, H_p(F, \mathbf{C})).$$

Date: December 2, 2001.

2000 Mathematics Subject Classification. 20F55.

Both authors acknowledge the support of NSERC Postdoctoral Fellowships.

This is a refinement of the usual zeta function of the monodromy of the Milnor fibre, which Denef and Loeser [8] have calculated for all reflection groups. Their technique uses Springer's theory of regular elements [19]: $g \in G$ is called a *regular element* of G iff it has an eigenvector that is not contained in any reflecting hyperplane.

In particular, Springer [19] has shown that the centralizer of a (noncentral) regular element in G acts as a reflection group on a (proper) subspace of V . Using this idea and its elaboration in [8, 14], we find a recursive formula for the Euler characteristic χ_Γ (Theorem 3.13).

For a fixed reflection group G , let $M_G = M = V - \cup_{H \in \mathcal{A}} H$ be the hyperplane complement, and U its image in $\mathbf{P}(V)$. These spaces have been studied extensively in the context of hyperplane arrangement theory; see, for example, [13]. On the other hand, let \mathcal{E} denote the set of all maximal eigenspaces E of elements of G for which $E \subsetneq V$. Let

$$(1.2) \quad M^\circ = V - \bigcup_{E \in \mathcal{E}} E,$$

and U° the image of M° in $\mathbf{P}(V)$. U° is the complement of a projective subspace arrangement, in the sense of Björner [1], and to the authors' knowledge has not been studied directly before.

$G/Z(G)$ acts freely on U° . We find a formula for the Euler characteristic of the orbit space in terms of degrees, codegrees, and regular numbers (4.22), and we calculate it for each irreducible G (Theorem 3.15). We show that this determines χ_Γ for each G (Theorem 3.13).

2. SPRINGER'S THEORY OF REGULAR ELEMENTS

In this section, we recall the theory of regular elements and set up our notation. We refer to [16] for background on reflection groups and hyperplane arrangements and to [19] for background on the theory of regular elements.

Let $V = \mathbf{C}^\ell$ and let G be a finite reflection group acting on V . We will denote by $\mathbf{C}[V]$ the algebra of polynomial functions on V . The *degrees* d_1, \dots, d_ℓ of G are the degrees of any set of homogeneous polynomials which generate the G -invariant polynomial ring $\mathbf{C}[V]^G$. The order of G and of its centre $Z(G)$ are determined in terms of its degrees:

$$(2.3) \quad |G| = \prod_{i=1}^{\ell} d_i \quad |Z(G)| = \gcd \{d_i\}$$

A vector $v \in V$ is called *regular* if it is not contained in a reflection hyperplane of G . An element $g \in G$ is called *regular* if it has a regular eigenvector. Let $g \in G$ be regular of order d . Let v be a regular eigenvector with corresponding eigenvalue ξ and let $V(g, \xi)$ denote the ξ -eigenspace of g . We will refer to (g, ξ) as a *regular (d -)pair*.

With this notation, we have:

Theorem 2.4 (Springer [19]).

- (a) *The root of unity ξ has order d .*
- (b) *$V(g, \xi)$ has dimension $a(d) = |\{i : d|d_i\}|$.*
- (c) *The centralizer $C_G(g)$ is a reflection group in $V(g, \xi)$ whose degrees are $\{d_i : d|d_i\}$ and whose order is $\prod_{d|d_i} d_i$.*

The orders of the regular elements of G are called the *regular numbers* of G . Let \mathcal{R} denote the poset of regular numbers, ordered by divisibility.

The group $G \leq GL(V)$ also acts naturally on the algebra of polynomial vector fields on V , $\mathbf{C}[V] \otimes V$. The module $(\mathbf{C}[V] \otimes V)^G$ is free over $\mathbf{C}[V]^G$. Following [2], the *codegrees* d_1^*, \dots, d_ℓ^* are defined to be the degrees of a homogeneous basis, with the convention that derivations have degree -1 . By a theorem of Orlik and Solomon [18]

$$(2.5) \quad \sum_{i=1}^l \dim H^i(U, \mathbf{C}) t^i = \prod_{i=2}^\ell (1 + (d_i^* + 1)t)$$

Using a case-based argument, Denef and Loeser [8, Theorem 2.8] proved that, for a regular d -pair (g, ξ) , the codegrees of $C_G(g)$ acting on $V(g, \xi)$ are

$$(2.6) \quad \{d_i^* : d|d_i^*\}$$

Lehrer and Springer [14, Theorem C] later reproved this result in a case-free way.

3. EULER CHARACTERISTICS

Following [2], for each G -orbit of hyperplanes $\mathcal{C} \in \mathcal{A}/G$, set

$$\delta_{\mathcal{C}} = \prod_{H \in \mathcal{C}} \alpha_H^{e_{\mathcal{C}}},$$

where $e_{\mathcal{C}}$ is defined to be the common value of e_H for all $H \in \mathcal{C}$. noting that e_H is constant for all $H \in \mathcal{C}$. Consider any homogeneous, G -invariant polynomial $f \in \mathbf{C}[V]^G$ with zero locus equal to $\bigcup_{H \in \mathcal{A}} H$. Then f has the form

$$(3.7) \quad f = \prod_{\mathcal{C} \in \mathcal{A}/G} \delta_{\mathcal{C}}^{a_{\mathcal{C}}}$$

for some positive integers $a_{\mathcal{C}}$; in particular, the discriminant is obtained by choosing all $a_{\mathcal{C}} = 1$. We shall call such f *unreduced discriminant polynomials*.

Denote the degree of f as above by m . Then $f(gv) = f(\zeta v) = \zeta^m f(v)$ for any regular element g with eigenvalue v , so the order of ζ must divide m . That is, all regular numbers $d \in \mathcal{R}$ divide m .

Let $F = f^{-1}(1)$. Let $P = \pi_1(M, 1)$, the pure Artin braid group corresponding to the group G . Since F is homotopy-equivalent to an infinite cyclic cover of M , we have $H_*(F, \mathbf{C}) \cong H_*(M, \mathbf{C}[t, t^{-1}])$, where $\mathbf{C}[t, t^{-1}] \cong \mathbf{C} \uparrow_{\pi_1(F)}^P$ as a P -module; see [9, Section 2.1] and [10]. Explicitly, P has a set of generators $\{\gamma_H : H \in \mathcal{A}\}$ for which γ_H acts by multiplication by $t^{a_{\mathcal{C}} e_{\mathcal{C}}}$.

The complement M is known to be a $K(P, 1)$ space for all irreducible reflection groups with the possible exception of $G_{24}, G_{27}, G_{29}, G_{31}, G_{33}$, and G_{34} ; see [2, (2.11)] for references. Thus we have $H_*(F, \mathbf{C}) \cong H_*(P, \mathbf{C}[t, t^{-1}])$ except perhaps in these cases. Let $B = \pi_1(M/G, 1)$, the braid group. Then $H_*(P, \mathbf{C}[t^{\pm 1}])^G \cong H_*(B, \mathbf{C}[t^{\pm 1}])$ (over \mathbf{C}). For all real reflection groups, this is computed explicitly in [7].

The Lefschetz zeta function of F is defined to be

$$Z(F) = \prod_{p \geq 0} \det(1 - h_* t|_{H_p(F, \mathbf{C})})^{(-1)^p},$$

where h^* denotes a preferred generator of C_m acting in homology. Since a complex representation of C_m is determined by the characteristic polynomial of a generator of the group, the zeta function can be seen as the restriction to C_m of the Euler characteristic $\chi_\Gamma(F)$ defined in (1.1), written multiplicatively. We will identify $C_m = \langle h^* \rangle$ with the cyclic group of m elements in \mathbf{C}^* . For convenience, we will take the convention that $\alpha \in C_m$ acts on F by multiplication by α^{-1} .

For a reflection group G and integers $d|m$, define

$$I_d(G) = I_d = 1 \uparrow_{C_d}^{G \times C_m}$$

whenever C_d is a cyclic subgroup of order d generated by a regular pair (g, ζ) of order d .

Lemma 3.8. *The cyclic groups generated by any two regular pairs of the same order are conjugate in Γ . In particular, the definition of $I_d(G)$ above does not depend on the choice of regular pair (g, ζ) . The normalizer in Γ of the cyclic subgroup generated by $(g, \xi) \in \Gamma$ is $C_G(g) \times C_m$.*

Proof. Let (g, ζ) and (g', ξ) be two regular pairs of order d , generating cyclic groups K and K' , respectively. Since any two primitive roots of unity of the same order generate the same cyclic subgroup of \mathbf{C}^* , $\xi = \zeta^k$ for some k . Then (g^k, ξ) is also a regular pair of order d , so g^k and g' are conjugate, by [19, 4.2]. It follows that the subgroups K and K' are conjugate. $I_d(G)$ is well defined since the permutation characters induced from K and K' are the same [6, 10.12]. Since $(h, \alpha) \in N_\Gamma((g, \xi))$ iff $(hgh^{-1}, \alpha) = (g^k, \xi^k)$ for some k iff $h \in C_G(g)$, $\alpha \in C_m$, we have that $N_\Gamma(\langle (g, \xi) \rangle) = C_G(g) \times C_m$. ■

The following theorem appeared independently as [8, Theorem 2.5] and [13, Corollary 5.8].

Lemma 3.9. *If (g, ζ) is a regular pair and V is the ζ -eigenspace of g , then the centralizer $C(g)$ acts as a reflection group on V . Its reflecting hyperplanes are $\{H \cap V : H \in \mathcal{A}\}$.*

Definition 3.10. For a regular d -pair (g, ζ) , we define $U(g, \zeta)$ as the projective hyperplane complement for $C_G(g)$ acting on $V(g, \zeta)$ where (g, ζ) is a regular d -pair.

From Lemma 3.8, $C_G(g)$ is conjugate to $C_G(g')$ if (g, ζ) and (g', ξ) are both regular d -pairs. This means that they are isomorphic as reflection groups, and we will refer to them, up to isomorphism, as $G(d)$, as in [14].

By the lemma above, then, $U(g, \zeta)$ and $U(g', \xi)$ are diffeomorphic; we will refer to them as $U_G(d)$.

Definition 3.11. Define a poset $\mathcal{D} = \mathcal{D}_G$ by

$$\mathcal{D} = \{d : d = |Z(C_G(g))| \text{ for a regular element } g\},$$

ordered by divisibility. That is, \mathcal{D} is the set of orders of regular elements g that are maximal with respect to the property of having a given centralizer. For elements $d \in \mathcal{R}_G$, define $[d]$ to be the least multiple of d in \mathcal{D} .

Note that $\{G(d) : d \in \mathcal{D}\}$ forms a complete set of representatives of the isomorphism classes of centralizers of regular elements.

Recall $M^\circ \subseteq M$ from (1.2), and $U^\circ \subseteq U$. By construction,

Proposition 3.12. $G/Z(G)$ acts freely on U° .

We can now state our main result.

Theorem 3.13. Let G be a reflection group, f an unreduced discriminant polynomial (3.7) of degree m , and F its Milnor fibre. Then

$$\chi_\Gamma(F) = \sum_{d \in \mathcal{D}} a_d I_d,$$

where the integers a_d are given by

$$(3.14) \quad a_d = \chi(U(d)^\circ / G(d))$$

A case analysis gives a more refined description. First, $\chi_\Gamma(F)$ is zero unless G is irreducible, since $\chi(U) = 0$ in this case: this appears first in the language of matroids in [5]. Denef and Loeser [8, 2.9] show that the centralizers of regular elements in irreducible G are themselves irreducible. With this in mind, it is enough to calculate $a_z = \chi(U^\circ / G)$, where $z = |Z(G)|$, for each irreducible G . We obtain:

Theorem 3.15. For an irreducible reflection group G of rank n , $\chi(U^\circ / G) = (-1)^{n-1}$ if G is in the list below. Otherwise, $\chi(U^\circ / G) = 0$.


- (a) Irreducibles of rank ≤ 2 ;
- (b) Irreducibles of the form $G(d\ell, e\ell, \ell)$, except $G(3, 3, 3)$;
- (c) $G(d\ell, e\ell, 2\ell)$ where $e\ell$ is odd;
- (d) Exceptionals G_{29} and G_{34} .

Corollary 3.16. For a Milnor fibre of a given reflection group as above, χ_Γ is a linear combination of at most six permutation characters I_d , with coefficients ± 1 .

The value of χ_Γ for each irreducible reflection group is tabulated in Section 6.

We also observe empirically that, like the zeta function, χ_Γ continues to be a braid diagram invariant, in the sense of Broué, Malle, and Rouquier [2].

Example 3.17. Let G be the irreducible reflection group of type E_8 . The degrees are $[2, 8, 12, 14, 18, 20, 24, 30]$ and the poset \mathcal{D} is:



By the Shephard-Todd classification, the centralizers of elements of order 4 and 6 are, respectively, G_{31} and G_{32} . Applying Theorems 3.13 and 3.15 shows that

$$\chi_{E_8 \times C_m} = I_{30} + I_{24} + I_{20} - I_{12} - I_{10} - I_8.$$

The rest of this paper is as follows. Theorem 3.13 is proven in Section 4. Theorem 3.15 is proven in Section 6. We are unable to provide a case-free proof of Theorem 3.15 in general, although in Section 5 we do so for rank 2 irreducibles.

4. PROOF OF THEOREM 3.13

Throughout this section, fix a particular reflection group G acting on $V = \mathbf{C}^\ell$. We will use the notation of Section 3.

For a regular element $g \in G$ of order d , let $i_G(d)$ be the order of its centralizer, and let $u_G(d)$ be the Euler characteristic of $U(d)$. Then $i(d) = \prod_{d|d_i} d_i$ (Theorem 2.4). By evaluating (2.5) at $t = -1$ and using (2.6), we see $u(d) = \prod_{d|d_i^*, i > 1} (-d_i^*)$.

If (g, ζ) is a regular pair of order d , then by [19, 3.4], there is a regular pair (h, ζ) of order $\lceil d \rceil$ for which $C(g) = C(h)$ (and, hence, $U(g, \zeta) = U(h, \zeta)$). It follows that $i(d) = (\lceil d \rceil)$ and $u(d) = u(\lceil d \rceil)$.

For $(g, \zeta) \in G \times C_m$, it follows from a refinement of the Hopf-Lefschetz fixed point theorem [3] that

$$(4.18) \quad \begin{aligned} \chi_\Gamma(F)(g, \zeta) &= \chi(F^{(g, \zeta)}) = \deg(\delta) \cdot \chi(U(g, \zeta)) \\ &= \begin{cases} m \cdot u(\lceil d \rceil) & \text{if } (g, \zeta) \text{ is regular of order } d; \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Lemma 4.19. *There exist rationals $\{a_d : d \in \mathcal{D}\}$ for which*

$$(4.20) \quad \chi_\Gamma = \sum_{d \in \mathcal{D}} a_d I_d.$$

Proof. We claim that, if I_d is induced from a regular pair (g, ζ) of order d , then

$$(4.21) \quad I_d(h, \xi) = \begin{cases} \frac{m}{d} |C(h)| & \text{if } (h, \xi) \text{ is regular of order } d', d'|d; \\ 0 & \text{otherwise.} \end{cases}$$

This follows by directly evaluating the induced character and using Lemma 3.8: $I_d(h, \xi)$ is nonzero only if (h, ξ) is conjugate to a power of (g, ζ) , the generator of C_d , which is equivalent to (h, ξ) being a regular pair of an order dividing that of g .

Define an equivalence relation \sim on $G \times C_m$ by setting $(g, \zeta) \sim (h, \xi)$ iff either: neither is a regular pair; or, both are regular pairs, and $[g] = [h]$.

The equivalence classes of \sim are unions of conjugacy classes. Moreover, (4.21) shows that each I_d is constant on classes of \sim .

The characters I_d span the \mathbf{Q} -vector space of functions that are both constant on \sim classes and zero on the nonregular equivalence class, since their values on the regular pairs form a triangular matrix, by (4.21).

By (4.18), χ_Γ is such a class function, which completes the proof. ■

4.1. coefficients a_d from (co)degrees. By evaluating (4.20) on a regular pair of order $d \in \mathcal{D}$, we obtain

$$m \cdot u(d) = \sum_{k \in \mathcal{D}: d|k} a_k \frac{m}{k} i(d),$$

whence by Möbius inversion,

$$\begin{aligned} (4.22) \quad a_d &= d \sum_{d|k} \mu(d, k) \frac{u(k)}{i(k)}, \\ &= d \sum_{d|k} \mu(d, k) \prod_{\substack{i:d|d_i^* \\ d_i^* \neq 0}} d_i^* \prod_{i:d|d_i} d_i^{-1} \end{aligned}$$

where μ is the Möbius function of the poset \mathcal{D} .

Remark 4.23. Any rational character of a finite group can be expressed as a rational linear combination of induced permutation characters from cyclic subgroups by the Artin-Brauer induction theorem [6, 15.2]. The coefficients can be determined and are non-zero only if the character is non-zero on that cyclic subgroup. In our situation, however, the cyclic subgroups $C_d, d \in \mathcal{D}$ are representatives of isomorphism classes (hence a subset of the set) of cyclic subgroups of Γ with $\chi_\Gamma(F)$ non-zero. Our argument above is then a direct way to compute the coefficients $a_d, d \in \mathcal{D}$ for our special case.

4.2. induction from a centralizer subgroup. Given a regular element $g_0 \in G$ of order $e \in \mathcal{D}$, let $H = C_G(g_0)$. From [19, 4.2], and Definition 3.11 we observe:

Lemma 4.24. The maximal regular numbers of H are $\mathcal{D}_H = \{d \in \mathcal{D}_G : e|d\}$.

By Lemma 4.19,

$$\begin{aligned} \chi_{H \times C_m} &= \sum_{d \in \mathcal{D}_H} a'_d I_d(H), \quad \text{and} \\ \chi_{G \times C_m} &= \sum_{d \in \mathcal{D}} a_d I_d \end{aligned}$$

for some coefficients $\{a'_d\}$, and $\{a_d\}$.

Theorem 4.25. *If G and H are as above, and $m'|m$, then*

$$\chi_{G \times C_m} = \chi_{H \times C'_m} \uparrow_{H \times C'_{m'}}^{G \times C_m} + \sum_{d \in \mathcal{D}: e \nmid d} a_d I_d.$$

Consequently, in the notation above, $a_d = a'_d$ for all $d \in \mathcal{D}_H$.

Proof of theorem. We need to show that the values a_d given by equation (4.22) for $d \in \mathcal{D}_H$ are the same in H as they are in G . Specifically, we need to show for multiples k of e , that $i_G(k) = i_H(k)$, and that $u_G(k) = u_H(k)$.

So suppose that $g \in H$ is a regular element of order k , where $e|k$. Then g_0 is conjugate to a power of g , so $C_G(g) \subseteq C_G(g_0) = H$; that is, $C_G(g) = C_H(g)$, so $i_G(k) = i_H(k)$.

Now assume, without loss of generality, that $g = g_0^r$ for some r , and choose ζ so that (g, ζ) is a regular pair. Then the ζ eigenspace of g in V_G is contained in V_H , so $U_G(g, \zeta) = U_H(g, \zeta)$ (Lemma 3.9.) Then $u_G(k) = u_H(k)$ as well. \blacksquare

4.3. interpretation of coefficients a_d . For $d \in \mathcal{D}$, let $F_d = F \cap V_d$, where

$$V_d = \bigcap_{d \nmid d_i} f_i^{-1}(0),$$

the variety of eigenvectors of elements of g having eigenvalue a primitive d th root of unity; see [19, 3.2]. In particular, $F_z = F$, where z is the order of the centre of G . Let

$$F_d^\circ = F_d - \bigcup_{\substack{d|d' \\ d' \neq d}} F_{d'}.$$

The following lemma is standard.

Lemma 4.26. *Suppose $X \subseteq M$ is a smooth embedding of complex manifolds without boundary and X is closed in M . Putting $U = M - X$, we have*

$$\chi(X) + \chi(U) = \chi(M).$$

Furthermore, if a finite group G acts on M so that $GX = X$, we have

$$\chi_G(X) + \chi_G(U) = \chi_G(M).$$

Proposition 4.27. *For any $d \in \mathcal{D}$,*

$$\chi_{G \times C_m}(F_d) = \sum_{\substack{d' \in \mathcal{D} \\ d|d'}} \chi_{G \times C_m}(F_{d'}^\circ).$$

Proof. Use induction on the dimension of F_d . If minimal, $F_d = F_d^\circ$ and there is nothing to prove. Otherwise, we have

$$\begin{aligned} F_d &= F_d^\circ \cup \bigcup_{\substack{d|d' \\ d' \neq d}} F_{d'}^\circ \\ &= F_d^\circ \cup \bigcup_{d' \succ d} F_{d'}^\circ \end{aligned}$$

where “ \succ ” denotes the covering relation in the poset \mathcal{D} , and all of the unions are disjoint. The proof is completed by applying Lemma 4.26 and the induction hypothesis to the second equality. ■

To complete the proof of Theorem 3.13, it remains to show that the coefficients a_d given by Lemma 4.19 satisfy $a_d = \chi(U(g, \zeta)^\circ / C_G(g))$ for regular pairs (g, ζ) of order $d \in \mathcal{D}$.

Let $z = |Z(G)|$. From Proposition 3.12, the quotient map $F_z^\circ \rightarrow U^\circ / G$ is a covering with deck transformation group $G \times C_m / \langle (g, \zeta) \rangle$, where (g, ζ) is any regular z -pair. It follows that

$$\begin{aligned} \chi_{G \times C_m}(F^\circ) &= \chi(U^\circ / G) \cdot 1 \uparrow_{C_z}^{G \times C_m}, \\ &= \chi(U^\circ / G) \cdot I_z. \end{aligned}$$

By Theorem 4.25, then, for each $d \in \mathcal{D}$,

$$\chi_\Gamma(F_d^\circ) = \chi(U(d)^\circ / G(d))I_d,$$

Now, using Proposition 4.27 with $d = z$, we have

$$\begin{aligned} \chi_\Gamma(F) &= \sum_{d \in \mathcal{D}} \chi_\Gamma(F_d^\circ) \\ &= \sum_{d \in \mathcal{D}} a_d I_d; \end{aligned}$$

equating the coefficient of I_d for each d gives the characterization of the values a_d that we claimed.

5. THE RANK TWO CASE

For finite groups acting on \mathbf{C}^2 , a stronger version of Theorem 3.15(a) is obtained from the theory of du Val singularities, for which we refer to [11].

Theorem 5.28. *Let G be an irreducible finite subgroup of $U_2(\mathbf{C})$. Then*

$$\chi(U_G^\circ / G) = -1.$$

Proof. Note that a finite subgroup G of $GL_\ell(\mathbf{C})$ can always be embedded in $U_\ell(\mathbf{C})$. Let $Z := Z(U_\ell(\mathbf{C})) = \{\alpha I : \alpha\bar{\alpha} = 1\}$. Let $H := (G \cdot Z) \cap SU_\ell(\mathbf{C})$; then $Z(H) = H \cap SU_\ell(\mathbf{C})$. It is easy to check that the map $G/Z(G) \rightarrow H/Z(H)$ defined by $gZ(G) \mapsto g(\det(g)^{1/\ell})Z(H)$ is an isomorphism; therefore G and H are both central extensions of the same group \overline{H} . Then G and H act on \mathbf{C}^ℓ and \overline{H} acts on $\mathbf{P}^{\ell-1}$. Proper eigenspaces for the action of G

on \mathbf{C}^ℓ are the same as those for H . Moreover, if v is an eigenvector for an element $h \in H$, its image $[v] \in \mathbf{P}^{\ell-1}$ is a fixed point of $hZ(H) \in \overline{H}$.

Now, let $\ell = 2$. Note that for $G \leq U_2(\mathbf{C})$, there exists a reflection group $G' \leq U_2(\mathbf{C})$ with $\overline{H} = G/Z(G) \cong G'/Z(G')$. Then $\mathbf{P}^1/\overline{H} \cong \mathbf{P}^1/G \cong \mathbf{P}^1/G' \cong \mathbf{P}^1$ where the last isomorphism follows from the Shephard-Todd-Chevalley theorem.

Since $PSL_2(\mathbf{C}) \cong SO_3$, \overline{H} is isomorphic to a finite subgroup of SO_3 . The finite subgroups of SO_3 are known: they are the groups of symmetry of the regular polyhedra: cyclic, dihedral, tetrahedral (A_4), octahedral (S_4) and icosahedral (A_5). Since G was assumed to act irreducibly, \overline{H} is not cyclic. For the remaining finite subgroups of SO_3 , Klein [12] showed that there are exactly three orbits of points in S^2 of points with nontrivial stabilizers (corresponding to vertices, barycenters of edges of the regular polyhedron, and barycenters of faces.) Since $\mathbf{P}^1/G \cong \mathbf{P}^1$, it follows that $U_G^\circ/G = \mathbf{P}^1 - \{p_0, p_1, p_2\}$, where the p_i 's are the 3 “special” orbits under the action of H on \mathbf{P}^1 . Thus $\chi(U_G^\circ/G) = 2 - 3 = -1$. ■

6. PROOF OF THEOREM 3.15

The Shephard-Todd classification of irreducible (complex) reflection groups consists of one infinite family and 34 exceptional groups labelled as G_4, \dots, G_{37} . The tables in Figures 1, 2, 3 give χ_Γ for the exceptional groups; these values are readily calculated from (4.22).

For any reflection group G , put $c(G) = \chi(U^\circ/G)$. From Theorem 3.13, this equals the coefficient of $I_{|Z(G)|}$ in χ_Γ . By Theorem 4.25, for $d \in \mathcal{D}$, the coefficient a_d of I_d is $c(G(d))$.

Theorem 3.15 claims that for exceptional irreducible G , $c(G) = 0$ unless G has rank $\ell = 2$ or G is one of G_{29} or G_{34} , in which case $c(G) = (-1)^{\ell-1}$. Our proof in ranks $\ell > 2$ is by inspection, after having computed χ_Γ for each group.

The rest of this section is devoted to proving Theorem 3.15 for the infinite family of reflection groups $G(r, p, \ell)$.

6.1. (co)degrees and regular numbers. For $r, p \in \mathbf{N}$ with $p|r$ and rank $\ell \geq 2$, $G(r, p, \ell)$ is a group of order $r^\ell/p\ell!$. It is the semidirect product of the symmetric group S_ℓ acting by permutations on the standard basis $\{e_i : 1 \leq i \leq \ell\}$, and the group of diagonal maps $e_i \mapsto \theta_i e_i$, where $\theta_i^r = 1$ and $(\theta_1 \cdots \theta_\ell)^q = 1$ where $q = r/p$. The group acts irreducibly on \mathbf{C}^ℓ iff $r > 1$ and $(r, p, \ell) \neq (2, 2, 2)$. $G(r, 1, \ell)$ is the full monomial group $C_r^\ell \rtimes S_\ell$ and that the Weyl groups $A_{\ell-1}, B_\ell, D_\ell, G_2$ and the dihedral groups $I_2(\ell)$ equal respectively $G(1, 1, \ell), G(2, 1, \ell), G(2, 2, \ell), G(6, 6, 2)$ and $G(\ell, \ell, 2)$. The codegrees are calculated in [17], and the regular numbers appear in [4]. The degrees of $G(r, p, \ell)$ are

$$(6.29) \quad \begin{cases} r, 2r, \dots, (\ell-1)r, \ell q, & p|r, r > 1 \\ 2, 3, \dots, (\ell-1), \ell, & p = r = 1. \end{cases}$$

the order of the center z is $q \gcd(r, \ell)$. The codegrees are

$$(6.30) \quad \begin{cases} 0, r, 2r, \dots, (\ell - 1)r, & p < r \\ 0, r, 2r, \dots, (\ell - 2)r, (\ell - 1)r - \ell & p = r > 1, \ell > 1 \\ 0, 1, 2, \dots, (\ell - 2), & p = r = 1, \ell > 2 \end{cases}$$

Note that the adjustments in the degrees and codegrees of $G(1, 1, \ell) = S_\ell$ are made so that S_ℓ acts irreducibly.

Remark 6.31. Since the degrees and codegrees of $G(d)$ are those of G which are divisible by d , we see for $G = G(r, p, \ell)$ and $e = (\gcd(d, r))^{-1}d$, the degrees of $G(d)$ are

$$(6.32) \quad \begin{cases} er, 2er, \dots, \lfloor \frac{\ell-1}{e} \rfloor er, & d \nmid \ell q \\ er, 2er, \dots, \lfloor \frac{\ell-1}{e} \rfloor er, \ell q, & d \mid \ell q \\ 2, 3, \dots, \ell, & d = r = 1, \end{cases}$$

as noted in [14]. Lehrer and Springer prove indirectly in [14, 5.2] that $G(d) = G(r', p', \ell')$ for $G = G(r, p, \ell)$ and $d \in \mathcal{R}$. In the proposition below we make the determination of $G(d)$ explicit to help in our computation of $c(G)$.

Proposition 6.33.

- (a) For $G = G(r, p, \ell)$, $q = r/p > 1$, we have $\mathcal{R} = \{d : d \mid \ell q\}$ and $\mathcal{D} = \{kq : t \mid k \mid \ell\}$ where $t = \gcd(p, \ell)$. Then for $kq \in \mathcal{D}$, $G(kq) = G(kr/t, p, \ell t/k)$.
- (b) For $G = G(r, r, \ell)$, $\ell > 1$, we have $\mathcal{R} = \{d : d \mid \ell q\} \cup \{d : d \mid (\ell - 1)r\}$ and

$$\mathcal{D} = \{kr : k \mid \ell - 1\} \cup \{d : z \mid d \mid \ell\}$$

where $z = \gcd(r, \ell)$. For $k \mid \ell - 1$, $kr \neq z$, we have $G(kr) = G(kr, 1, (\ell - 1)/k)$ and for $z \mid d \mid \ell$, we have $G(d) = G(dr/z, r, \ell z/d)$.

Proof. In each case, let $z = |Z(G)|$, equal to the greatest common divisor of the degrees. In the first case, $z = \gcd(pq, \ell q) = tq$, where $t = \gcd(p, \ell)$. In the second case, this is just $z = \gcd(r, \ell)$.

Case (a): For $G = G(r, p, \ell)$ with $q > 1$, the only maximal regular degree of G is ℓq [4, 2.11]. Thus the set of regular numbers are $\mathcal{R} = \{d : d \mid \ell q\}$. If $d \in \mathcal{D}$, then d is a gcd of a subset of the degrees. So $qt \mid d$, since $z = qt$ is the gcd of all the degrees. To show that

$$\mathcal{D} = \{kq : \gcd(p, \ell) \mid k \mid \ell\}$$

we need to show that $G(d)$ are distinct for distinct d in this set. Observe that if $\gcd(p, \ell) \mid k \mid \ell$, then $\gcd(p, k) = \gcd(p, \ell)$. But as is implicitly shown in the proof of [14, 5.2] we have for $t \mid k \mid \ell$,

$$G(kq) = G(kr/t, p, \ell t/k)$$

So we have shown that \mathcal{D} and $G(d), d \in \mathcal{D}$ are described as in the claim.

Case (b): For $G = G(r, r, \ell)$, with $\ell > 1$, the maximal regular degrees are $(\ell - 1)r$ and ℓ [4, 2.11]. So the set of regular numbers is

$$\mathcal{R} = \{d : d|\ell\} \cup \{d : d|(\ell - 1)r\}$$

For $d \in \mathcal{D}$, $z = \gcd(r, \ell)$, and $z|d$ since d is the gcd of a subset of the degrees. If $d \in \mathcal{D}$ does not divide ℓ , then $d|(\ell - 1)r$ and d is a gcd of a subset of $\{r, \dots, (\ell - 1)r\}$ so that $r|d$. We have shown that $\mathcal{D} \subseteq T_1 \cup T_2$, where

$$(6.34) \quad T_1 = \{d : z|d|\ell\} \quad \text{and} \quad T_2 = \{kr : k|\ell - 1\}.$$

Note that these sets intersect iff $\gcd(r, \ell) = r$, iff $r|\ell$. To show the inclusion is an equality, we have to show that $G(d)$ are distinct for distinct $d \in T_1 \cup T_2$. If $d \in T_1$, we have $\gcd(r, d) = \gcd(r, \ell) = z$, so that by the proof of [14, 5.2], $G(d) = G(dr/z, r, \ell z/d)$.

On the other hand, if $d = kr$ where $k|\ell - 1$ but $d \neq z$, we have by the proof of [14, Prop. 5.2] that $G(kr) = G(kr, 1, (\ell - 1)/k)$. These groups have distinct parameters for each $d \in T_1 \cup T_2$, so $\mathcal{D} = T_1 \cup T_2$ as claimed. ■

6.2. Proof of Theorem 3.15 for $G(r, p, \ell)$. It remains to show that $c(G) = 0$ for irreducible $G = G(r, p, \ell)$, except for the parameters $(d\ell, e\ell, \ell) \neq (3, 3, 3)$, and for $(d\ell, e\ell, 2\ell)$, where $e\ell$ is odd. For these exceptions, we show $c(G) = (-1)^{\ell-1}$. It will be convenient to let $S(m, k) = 1/m \sum_{d|m} \mu(d)(-1)^{md/k-1}$, where μ is the (number-theoretic) Möbius function on \mathbf{N} .

Lemma 6.35.

$$S(m, k) = \begin{cases} (-1)^{km-1}, & m = 1, \text{ or } m = 2 \text{ and } k \text{ odd} \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Recall that $\sum_{d|m} \mu(d) = \delta_{m,1}$. Consider the case where k is odd and m even. Write $m = 2^s t$ for t odd. Then

$$\begin{aligned} S(m, k) &= \frac{1}{m} \left(- \sum_{d|m, m/d \text{ even}} \mu(d) + \sum_{d|m, m/d \text{ odd}} \mu(d) \right) \\ &= \frac{1}{m} \left(- \sum_{d|m/2} \mu(d) - \sum_{d|m, 2^s|d} \mu(2^s) \mu\left(\frac{d}{2^s}\right) \right) \\ &= -\frac{2}{m} \delta_{m,2} = -\delta_{m,2}. \end{aligned}$$

The remaining cases are similar. ■

At the same time, we will calculate χ_Γ .

Proposition 6.36. *Let $\Gamma = G \times C_m$ and $m = \deg(\delta_G)$. Then*

(a) For $G = G(r, p, \ell)$ and $q = r/p > 1$,

$$\chi_\Gamma = \begin{cases} I_{\ell q}, & \ell \text{ odd}; \\ -I_{\ell q}, & \ell \text{ and } p \text{ even}; \\ I_{\ell q} - I_{q\ell/2}, & \ell \text{ even, } p \text{ odd.} \end{cases}$$

(b) For $G = G(r, r, \ell)$, and $\ell > 1$,

$$\chi_\Gamma = \begin{cases} I_{(\ell-1)r} - I_{r(\ell-1)/2} + I_\ell, & \ell \text{ odd}; \\ I_{(\ell-1)r} - I_\ell, & r, \ell \text{ even}; \\ I_{(\ell-1)r} + I_\ell - I_{\ell/2} & \ell \text{ even, } r \text{ odd.} \end{cases}$$

Proof. We will handle C_r as the special case $G(r, 1, 1)$ of (a) and S_ℓ as the special case $G(1, 1, \ell)$ of (b). Recall that $n = \text{rk}(G(r, p, \ell)) = \ell$ unless $r = p = 1$ when $n = \text{rk}(G(1, 1, \ell)) = \ell - 1$ since $G(1, 1, \ell) = S_\ell$ acts irreducibly on an $\ell - 1$ dimensional space.

(a) For $G = G(r, p, \ell)$, $p < r$, we have

$$\mathcal{D} = \{kq : t|k|\ell\}$$

where $t = \gcd(p, \ell)$ and for $t|k|\ell$, $G(kq) = G(kr/t, p, \ell t/k)$. So

$$\begin{aligned} c(G) &= a_{tq} = tq \sum_{tq|kq} \mu(tq, kq) \frac{u(kq)}{i(kq)} \\ &= \frac{tq}{\ell q} \sum_{t|k|\ell} \mu(k/t)(-1)^{\ell t/k-1} \\ &= S(\ell/t, t) \\ &= \begin{cases} (-1)^{\ell-1} & \ell = t \text{ or } \ell = 2t, t \text{ odd } (\ell|p \text{ or } \frac{\ell}{2}|p, p \text{ odd.}) \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

For $G(kq) = G(kr/t, p, \ell t/k)$ with $t|k|\ell$ we have

$$c(G(kq)) = a_{kq} = \begin{cases} (-1)^{\ell t/k-1}, & k = \ell \text{ or } k = \ell/2, p \text{ odd} \\ 0, & \text{otherwise.} \end{cases}$$

This shows that

$$\chi_\Gamma = \begin{cases} I_{\ell q}, & \ell \text{ odd}; \\ -I_{\ell q}, & p, \ell \text{ even}; \\ I_{\ell q} - I_{q\ell/2}, & \ell \text{ even, } p \text{ odd,} \end{cases}$$

as required.

(b) For $G = G(r, r, \ell)$, $\ell > 1$, by Proposition 6.33, $\mathcal{D} = T_1 \cup T_2$, defined in (6.34). For $d \in T_1$, we have $G(d) = G(dr/z, r, \ell z/d)$, and for $d \in T_2$, $G(d) = G(kr, 1, (\ell - 1)/k)$. Since $z = \gcd(r, \ell)$, we have $z \in T_1$, and

$z \in T_2$ if and only if $z = r$.

$$\begin{aligned}
c(G) &= a_z = z \sum_{d \in \mathcal{D}} \mu(z, d) \frac{u(d)}{i(d)} \\
&= z \frac{u(z)}{i(z)} + z \sum_{d \in T_1 - \{z\}} \mu(z, d) \frac{u(d)}{i(d)} + z \sum_{kr \in T_2 - \{z\}} \mu(z, kr) \frac{u(kr)}{i(kr)} \\
&= (-1)^{\ell-1} \frac{z((\ell-1)r - \ell)}{\ell(\ell-1)r} + \frac{z}{\ell} \sum_{\substack{1 \neq \frac{d}{z} \mid \frac{\ell}{z} \\ 1 \neq \frac{d}{z} \mid \frac{\ell}{z}}} \mu\left(\frac{d}{z}\right) (-1)^{\frac{\ell}{d/z} z - 1} + \\
&\quad + \frac{z}{(\ell-1)r} \sum_{k \mid \ell-1, kr \neq z} \mu(z, kr) (-1)^{(\ell-1)/k-1} \\
&= (-1)^{\ell-1} \frac{z}{(\ell-1)r} + S(\ell/z, z) + \frac{z}{(\ell-1)r} \sum_{k \mid \ell-1, kr \neq z} \mu(z, kr) (-1)^{(\ell-1)/k-1}
\end{aligned}$$

Note that if $z \neq r$ then $\mu(z, r) = -1$ and $\mu(z, kr) = 0$ for all $1 \neq k \mid \ell-1$ whereas if $z = r$ then $\mu(z, kr) = \mu(kr/z) = \mu(k)$. So we have

$$c(G) = S(\ell/z, z) + \delta_{zr} S(\ell-1, 1)$$

This means that $c(G) = (-1)^{n-1}$ where $n = \text{rk}(G)$ iff

$$\begin{cases} \ell \mid r, & (r, \ell) \neq (2, 2), (3, 3) \\ \frac{\ell}{2} \mid r & r \text{ odd} \\ (r, \ell) = (1, 3) & \end{cases}$$

Otherwise $c(G) = 0$. Note that $c(G(2, 2, 2)) = 0$ agrees with the statement since $G(2, 2, 2)$ is reducible. Also observe that if $G = G(r, p, \ell)$ is a rank 2 irreducible, we obtain $c(G) = -1$ as was predicted in Section 5. This includes the case of the rank 2 irreducible $G(1, 1, 3) = S_3$.

To compute χ_Γ , it remains only to find a_d , for $d \in \mathcal{D}$. If $d \in T_1$, by part (a),

$$a_d = c\left(G\left(\frac{dr}{z}, r, \frac{\ell z}{d}\right)\right) = \begin{cases} (-1)^{\frac{\ell z}{d}-1} & d = \ell, \text{ or } d = \ell/2, r \text{ odd;} \\ 0 & \text{otherwise.} \end{cases}$$

For $d \in T_2$,

$$a_d = c\left(G\left(kr, 1, \frac{\ell-1}{k}\right)\right) = \begin{cases} 1 & d = (\ell-1)r; \\ -1 & d = r(\ell-1)/2, \ell \text{ odd;} \\ 0 & \text{otherwise.} \end{cases}$$

The expression for χ_Γ in (b) follows. ■

#	χ_Γ	#	χ_Γ
3	I_r	13	$I_{12} - I_4$
4	$I_6 + I_4 - I_2$	14	$I_{24} - I_6$
5	$I_{12} - I_6$	15	$-I_{12}$
6	$I_{12} - I_4$	16	$I_{30} + I_{20} - I_{10}$
7	$-I_{12}$	17	$I_{60} - I_{20}$
8	$I_{12} + I_8 - I_4$	18	$I_{60} - I_{30}$
9	$I_{24} - I_8$	19	$-I_{60}$
10	$I_{24} - I_{12}$	20	$I_{30} + I_{12} - I_6$
11	$-I_{24}$	21	$I_{60} - I_{12}$
12	$I_8 + I_6 - I_2$	22	$I_{20} + I_{12} - I_4$

FIGURE 1. χ_Γ for G_3 and rank-2 exceptionals

#	χ_Γ	#	χ_Γ
23	$I_{10} + I_6$	28	$I_{12} + I_8 - I_6 - I_4$
24	$I_{14} + I_6$	29	$I_{20} - I_4$
25	$I_{12} + I_9 - I_6$	30	$I_{30} + I_{20} + I_{12} - I_{10} - I_6 - I_4$
26	I_{18}	31	$I_{24} + I_{20} - I_{12} - I_8$
27	I_{30}	32	$I_{30} + I_{24} - I_{12}$

FIGURE 2. χ_Γ for exceptionals of rank 3 and 4

#	rank	χ_Γ
33	5	$I_{18} + I_{10}$
34	6	$I_{42} - I_6$
35	6	$I_{12} + I_8 - I_6 - I_4$
36	7	$I_{18} + I_{14}$
37	8	$I_{30} + I_{24} + I_{20} - I_{12} - I_{10} - I_8$

FIGURE 3. χ_Γ for exceptionals of rank ≥ 5

7. ORBIFOLD EULER CHARACTERISTICS

The orbifold Euler characteristic of a space X under the action of a group G is defined to be

$$\sum_{[g]} \chi(X^g / C_G(g)),$$

where the sum is taken over all conjugacy classes of G .

The orbifold Euler characteristic of the Milnor fibre F under the action of a $G \times C_m$ can be expressed in terms of the integers $\{a_d\}$ from (3.14), and we include its calculation here as an example.

Lemma 7.37. *For any reflection group G ,*

$$\chi(U/G) = \sum_{d \in \mathcal{D}} a_d.$$

Proof. We have a disjoint union $U = \bigcup_{d \in \mathcal{D}} G \cdot U(d)^\circ$. The claim follows by Proposition 3.12, Lemma 4.26, and the definition $a_d = \chi(U(d)^\circ)/G(d)$. \blacksquare

Theorem 7.38. *Let G be a reflection group, f an unreduced discriminant polynomial of degree m , and F its Milnor fibre. The orbifold Euler characteristic of F with respect to $\Gamma = G \times C_m$ equals*

$$\sum_{d \in \mathcal{D}} da_d,$$

in the notation of Section 3.

Proof. For a regular pair (g, ζ) , we have $F^{(g, \zeta)}/C(g, \zeta) = U^g/C_G(g)$. Recall that if (g, ζ) is not a regular pair, then the set of fixed points is empty, so we need only consider a sum over conjugacy classes of regular pairs in $G \times C_m$.

Using Lemma 3.8, there are $\phi(d)$ conjugacy classes of regular pairs of order d , for each regular number d . So we have

$$\begin{aligned} \sum_{[(g, \zeta)]} \chi(F^{(g, \zeta)}/C(g, \zeta)) &= \sum_{d \in \mathcal{R}} \phi(d) \chi(U^g/C_G(g)) \\ &= \sum_{d \in \mathcal{R}} \phi(d) a_{\lceil d \rceil} \\ &= \sum_{d \in \mathcal{D}} \sum_{d' \mid d} \phi(d') a_d \\ &= \sum_{d \in \mathcal{D}} da_d, \end{aligned}$$

where the second equality follows from Lemma 7.37 together with Lemma 3.9. \blacksquare

Remark 7.39. *By way of comparison, the ordinary or orbifold Euler characteristics of F/Γ are equal to the image of χ_Γ under homomorphisms from the character ring of Γ to \mathbf{Z} that take I_d to 1, or to d , respectively.*

8. CONCLUDING REMARKS

This investigation leaves the obvious open question of whether Theorem 3.15 could be proven in a more conceptual way. Our proof depends on knowing (co)degrees and regular numbers for each group, which are not reflected in the simplicity of the statement.

We also note that $\Gamma = G \times C_m$ is not the most general group for which these calculations make sense. In general one should replace G by $N(G)$, the normalizer of G in $U(V)$, and C_m by $\Lambda = C_m \rtimes \text{Gal}(K_m/K)$, where

K is the splitting field for $N(G)$ and K_m is the extension of K containing all m th roots of unity. Note that $\text{Gal}(K_m/K)$ is a finite group of order dividing $\phi(m)$ which acts on \mathbf{C} by inflation to $\text{Gal}(\mathbf{C}/K)$, since K_m/K is a Galois extension. This action can be extended to a diagonal action on \mathbf{C}^n which stabilizes $F = \delta^{-1}(1)$ since the coefficients of δ lie in K . Note that $\text{Gal}(K_m/K) \cap C_m = 1$ and that $\text{Gal}(K_m/K)$ normalizes C_m . The actions of $\text{Gal}(K_m/K)$ and $N(G)$ on F commute by construction. It may be interesting to examine the Λ -module structure of the equivariant Euler characteristic χ_Λ . It is probable that an answer would involve Springer's twisted regular numbers.

REFERENCES

1. A. Björner, *Subspace arrangements*, First European Congress of Mathematics, Vol. I (Paris, 1992), Birkhäuser, Basel, 1994, pp. 321–370.
2. M. Broué, G. Malle, and R. Rouquier, *Complex reflection groups, braid groups, Hecke algebras*, J. Reine Angew. Math. **500** (1998), 127–190.
3. K. S. Brown, *Complete Euler characteristics and fixed-point theory*, J. Pure Appl. Algebra **24** (1982), no. 2, 103–121.
4. A. M. Cohen, *Finite complex reflection groups*, Ann. Sci. École Norm. Sup. (4) **9** (1976), no. 3, 379–436.
5. H. Crapo, *A higher invariant for matroids*, J. Comb. Th. **2** (1967), 406–417.
6. C. W. Curtis and I. Reiner, *Methods of representation theory. Vol. I*, John Wiley & Sons Inc., New York, 1990, With applications to finite groups and orders, Reprint of the 1981 original, A Wiley-Interscience Publication.
7. C. De Concini, C. Procesi, M. Salvetti, and F. Stumbo, *Arithmetic properties of the cohomology of Artin groups*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **28** (1999), no. 4, 695–717.
8. J. Denef and F. Loeser, *Regular elements and monodromy of discriminants of finite reflection groups*, Indag. Math. (N.S.) **6** (1995), no. 2, 129–143.
9. G. Denham, *Hanlon and Stanley's conjecture and the Milnor fibre of a braid arrangement*, J. Algebraic Combin. **11** (2000), 227–240.
10. G. Denham, *The combinatorial Laplacian of the Tutte complex*, J. Algebra **242** (2001), no. 1, 160–175.
11. A. H. Durfee, *Fifteen characterizations of rational double points and simple critical points*, Enseign. Math. (2) **25** (1979), no. 1-2, 131–163.
12. F. Klein, *Lectures on the icosahedron and the solution of equations of the fifth degree*, revised ed., Dover Publications Inc., New York, N.Y., 1956, Translated into English by George Gavin Morrice.
13. G. I. Lehrer, *Poincaré polynomials for unitary reflection groups*, Invent. Math. **120** (1995), no. 3, 411–425.
14. G. I. Lehrer and T. A. Springer, *Reflection subquotients of unitary reflection groups*, Canad. J. Math. **51** (1999), no. 6, 1175–1193, Dedicated to H. S. M. Coxeter on the occasion of his 90th birthday.
15. J. W. Milnor, *Singular points of complex hypersurfaces*, Princeton University Press, 1968.
16. P. Orlik and H. Terao, *Arrangements of hyperplanes*, Grundlehren der Mathematischen Wissenschaften, no. 300, Springer-Verlag, 1992.
17. P. Orlik and L. Solomon, *Unitary reflection groups and cohomology*, Invent. Math. **59** (1980), no. 1, 77–94.

18. ———, *Arrangements defined by unitary reflection groups*, Math. Ann. **261** (1982), no. 3, 339–357.
19. T. A. Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.

E-mail address: denham@noether.uoregon.edu

UNIVERSITY OF OREGON, EUGENE, OR, USA 97403

E-mail address: nlemire@uwo.ca

UNIVERSITY OF WESTERN ONTARIO, LONDON, ONTARIO, CANADA N6A 5B7